



Differences Between Healthy and *Ganoderma boninense* Infected Oil Palm Seedlings Using Spectral Reflectance of Young Leaf Data

Aiman N.N. Azmi¹, Siti K. Bejo^{1,2,3*}, Mahirah Jahari¹, Farrah M. Muharam⁴ & Ian Yule⁵

¹Department of Biological and Agricultural Engineering, Faculty of Engineering, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

²Smart Farming Technology Research Centre, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

³Institute of Plantation Studies, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁴Department of Agriculture Technology, Faculty of Agriculture, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

⁵Institute of Agriculture and Environment, Massey University, Private Bag 11222, Palmerston North, New Zealand

*Corresponding author E-mail: skbejo@upm.edu.my

Received 19 September 2019; Accepted 10 July 2021; Available online 20 August 2021

Abstract: *Ganoderma boninense* (*G. boninense*) is the causal agent of basal stem rot (BSR) which significantly reduced the productivity of oil palm plantations in Southeast Asia. At early stage, the disease did not show any physical symptoms that could be seen with naked eyes resulted in detection difficulties. To date, there was no effective detection for this disease, and conventional methods such as manual and laboratory-based required trained specialists as well as time-consuming. Therefore, this study was conducted using hyperspectral remote sensing to investigate the differences in spectral reflectance of young leaf (frond one (F1) of healthy and *G. boninense* infected oil palm seedlings. The seedlings were inoculated with *G. boninense* pathogen at five months old. At five months after inoculation, 558 spectral signatures of F1 were extracted from acquired hyperspectral images. Noise removal was done to the extracted spectral signatures to remove outliers in the data. Then, the spectral signatures were averaged and plotted to observe the differences. Differences in reflectance of healthy and *G. boninense* infected seedlings were seen evidently in the near-infrared (NIR) region. Thus, this study showed evidence that F1 spectral reflectance has the ability to detect early stage of *G. boninense* infection at oil palm seedlings.

Keywords: *Ganoderma boninense*, BSR disease, Hyperspectral imaging, Oil palm seedlings, Spectral reflectance.

Introduction

Malaysia at present monopolises 28% of world palm oil production and currently the second world's largest exporter of palm oil after

Indonesia, with 33% of world exports. Many commercial plantation companies cultivated oil palms on a vast scale because of the high

yield and low production cost. Nevertheless, the production of oil palm in Malaysia has exposed to BSR disease, which caused by *G. boninense* pathogen. *G. boninense* is a white-rot fungus that able to degrade lignin component and damage the xylem pathway causing a vital problem in water and nutrient transportation to the aerial part of oil palm (Paterson, 2007; Shu'ud *et al.*, 2007). As a result, the symptoms appear similar to water and nutrient deficiency. Besides mature trees, oil palm seedlings are also prone to the infection where the symptoms develop earlier and dreadful (Sanderson, 2005).

The earliest visual symptoms of *G. boninense* infection at oil palm seedling is the development of fruiting body at bole, yellowing of older leaves as well as necrosis of basal frond, reflecting over 50% internal damage of stem base. However, early detection based on these symptoms is complicated and confounding because the fruiting body may appear unpredictably before or after leaf symptom development (Sariah *et al.*, 1994; Idris *et al.*, 2006). Consequently, healthy and mild infected seedlings are hard to differentiate and easily be overlooked. In addition, there is growth inhibition in terms of production of new leaves, height and girth as the infection progresses, due to low photosynthesis rate. Besides the visual symptoms, *G. boninense* infection can also be identified from brown discolouration of the inner longitudinal section of bole. Nevertheless, this type of detection is unfeasible for commercial plantation as the process is tedious, labour extensive and might lead to tree destruction.

Various techniques have been developed to detect BSR disease at early stage namely Ganoderma Selective Medium (GSM) (Darus & Seman, 1993), polymerase chain reaction-DNA (PCR-DNA) technique (Idris *et al.*,

2003), enzyme-linked immunosorbent assay-polyclonal antibody (ELISA-PAb) (Idris & Rafidah, 2008), GanoSken Tomography (Idris *et al.*, 2010), Field Spectroscopy (Izzuddin *et al.*, 2013), Mid-infrared spectroscopy (Liaghat *et al.*, 2014b) and Terrestrial Laser Scanning (Khairunniza-Bejo & Vong, 2014; Azuan *et al.*, 2019; Husin *et al.*, 2020). However, limitations of these techniques are cost-time efficiency, impractical for large plantation area, and need experts to execute the procedures. Nowadays, the standard practice to detect *G. boninense* in a plantation is by visual examination, which is labour intensive, time-consuming and human dependent.

Different studies have shown the capability of hyperspectral imaging in detecting symptomatic and asymptomatic plant diseases. For example, Bravo *et al.* (2003) and Moshou *et al.* (2005) have utilised visible-NIR hyperspectral imaging for the early detection of yellow rust disease in winter wheat. Susič *et al.* (2018) and Lu *et al.* (2018) utilised hyperspectral technology to detect and distinguish different causes of tomato stress and different stages of multi-tomato leaf disease, respectively. Kumar *et al.* (2012) investigated the applicability of aerial hyperspectral imaging for detection of greening in a citrus plantation. Similarly, Shafri & Hamdan (2009), Shafri *et al.* (2012) and Izzuddin *et al.* (2015) used airborne hyperspectral imaging to detect BSR disease in oil palm plantation. Meanwhile, Liaghat *et al.* (2014a) and Ahmadi *et al.* (2017) collected spectral signatures from young leaves from matured oil palms using spectroradiometer and developed classification models to separate the healthy, mild, medium and severe *G. boninense* infected oil palms. For oil palm seedlings, Shafri *et al.* (2011) and Izzuddin *et al.* (2017) have utilised APOGEE spectroradiometer to collect reflectance

spectra of young leaves after six months of inoculation with *G. boninense* pathogen. However, the foliar symptom had appeared after two months of inoculation. The result showed that there were difficulties to distinguish the healthy and mild infected seedlings due to overlapped reflectance spectra.

As mentioned earlier, previous studies were unable to differentiate the healthy and mild *G. boninense* infected seedlings. According to Zwiggelaar (1998), spectral reflectance of plant was affected by physical structure, chemical composition, and spectral properties of leaves. Meanwhile, changes in pigment levels of leaves commonly associated with physiological reactions, environmental stresses and diseases reactions (Chang, 1998; Gamon & Surfus, 1999; Gitelson *et al.*, 2001, Gitelson *et al.*, 2002). Furthermore, there is a need to fill the knowledge gap in BSR detection, particularly in oil palm seedling with inoculation period of fewer than six months. Therefore, this study was conducted to focus on detection of BSR at young seedling with the age of five months old, prior to symptoms expression. The analysis was done based on reflectance spectra taken from frond 1 (F1) of healthy and *G. boninense* infected seedlings. Besides, this study also conducted to determine the capability of hyperspectral imaging to detect *G. boninense* infection in oil palm seedlings prior to symptoms expression.

Materials & Methods

Experimental design

The study was carried out in a controlled environment at Universiti Putra Malaysia (UPM) Transgenic Greenhouse (2°59'33.10"N, 101°43'19.16"E), Serdang, Malaysia. The samples consisted of 28 oil palm seedlings (commercial standard crosses of Dura × Pisifera, DxP) at four months old

obtained from Sime Darby Plantation, Banting, Malaysia. The seedlings were placed inside the greenhouse before inoculation for a month to acclimate the condition. The temperature inside the greenhouse was set at 27°C, referring to Kamil & Omar (2016). All seedlings were watered and fertilised regularly at the same rate.

The acclimatised five months old seedling was transplanted into a polybag containing 90% topsoil, 10% sand and 6 cm × 6 cm × 6 cm rubberwood block (RWB) that colonised with *G. boninense* pathogen. The roots of the seedlings were placed on top direct contact with the inoculated RWB (Fig. 1) and covered with soil; this method was called sitting technique (Naidu *et al.*, 2018). Seedlings transplanted with uninoculated RWB acted as the control in this study. Polymerase chain reaction (PCR) test was conducted using two of the inoculated (U) seedlings after two months of inoculation to confirm the *G. boninense* infection.

Hyperspectral image acquisition was conducted at five months after inoculation using FirefLEYE S185 snapshot camera (Cubert GmbH, Ulm, Germany) with a spectral range of 450 nm- 950 nm (125 bands) and spectral sampling of 4 nm. The camera was mounted horizontally on a custom tripod that positioned 2.6 m from the ground level (Fig. 2). The system was controlled by data acquisition software, Cube-Pilot supplemented by the manufacturer. An image of a seedling was taken at a time against a black background board on a sunny clear day from 11:00 a.m. to 2:00 p.m. local time to get a natural illumination.

The camera was calibrated with white and dark references before each image acquisition so that the integration time is almost the same, which could reduce the effects of illumination

and detector sensitivity. A dark reference was done by closing the lens while the white reference was carried out using a white rectangular board (99% light reflection) positioned flat and close to the lens. Each collected spectrum was calibrated as:

$$\text{Reflectance} = \frac{(\text{Image}-\text{Dark})}{(\text{White}-\text{Dark})} \quad (1)$$



Fig. (1): Sitting technique where RWB was placed in direct contact with the roots of an oil palm seedling.




Fig. (2): Illustration of hyperspectral image acquisition setup inside the greenhouse.

Data pre-processing

Cube-Pilot software was used to extract the spectral signatures of F1 from the hyperspectral images. Since F1 was at the top of the oil palm crown, it was clearly visible in the images (Fig. 3). An average of 20 spectral signatures was collected randomly from the right and left leaflets of F1 for each seedling.



Fig. (3) Spectral extraction from the right and left leaflets of frond 1 (F1).

Thereafter, outliers in the spectral signatures were removed using the box plot method. Box plot statistically examined the data in a graphical way to describe the distribution of the data. The outliers were categorised as the spectral signature that has infrequent values from the rest of the spectral signatures.

Data analysis

Reflectance analysis was popularly used in previous studies to analyse the spectral signatures of plants. Plant stress level resulted from abiotic or biotic factors usually declines the concentration of leaves chlorophyll (as well as carotenoids) which affect the spectral reflectance of leaves and provide the opportunity to determine the plant health status remotely (Zarco-Tejada *et al.*, 2004; Malenovský *et al.*, 2006).

The reflectance analysis was conducted by calculating the average spectral signatures of healthy and *G. boninense* infected seedlings. The averages were then plotted against reflectance to observe the patterns and differences between healthy and *G. boninense* infected seedlings were determined.

Results & Discussion

Fig. (4) shows the result of reflectance analysis of F1 for healthy (H) and *G. boninense* infected (U) seedlings. The U seedlings demonstrated low reflectance in the NIR range (750 – 950 nm) while the H produced high reflectance. The pattern produced by U seedlings in NIR range was normal for diseased plants as a result of xylem destruction, which thus caused chlorophyll pigments reduction and water deficiency.

Furthermore, NIR wavelength could penetrate deeper through leaf pigments compared to the visible wavelength; therefore, changes of reflectance in the NIR range were more evident compared to changes in the visible range during the stress period (Liaghat *et al.*, 2014a). The changes were caused by the rupture of mesophyll cell wall of leaves (Gausman, 1977; Chappelle *et al.*, 1992; Slaton *et al.*, 2001; Ahmadi *et al.*, 2017) resulting in lower reflectance and higher absorbance of NIR. In addition, the results obtained in this study have coincided with Liaghat *et al.* (2014a), where healthy oil palms reflected higher light than *G. boninense* infected oil palms, primarily in the NIR range.

In this analysis, the H seedlings reflected a slightly higher visible light than U seedlings, which was contrary with theory studied by other researchers. Healthy plants typically generated lower visible reflectance than diseased plants, particularly in the green

wavelengths which represent higher chlorophyll content of leaves. Nonetheless, Shafri *et al.* (2011) had obtained a similar outcome as this study in which healthy seedlings produced higher reflectance in the green wavelength compared to *G. boninense* infected seedlings. This reflectance pattern might therefore be a specific spectral signature for oil palm seedlings since each plant has a specific spectral signature in various spectral regions as claimed by Schmidt & Skidmore (2003).

In this study, there were no physical symptoms of *G. boninense* infection appeared at the seedlings after five months of inoculation; however, the PCR test had generated positive results which indicated the presence of the infection. The late appearance of visual symptoms may vary due to the age of seedlings during inoculation, as younger palms potentially be affected by the *G. boninense* pathogen quicker than matured palms because of a not well-developed rooting system (Rees *et al.*, 2009). Thus, the physical symptoms of younger seedlings will appear early compared to older seedlings. Besides, Shafri *et al.* (2011) claimed that the differences in disease infection were also due to the resistance of the oil palm seedlings towards *G. boninense* pathogen.

The results have successfully filled the gap on the study of BSR detection at the very young age of seedlings (less than six months old) which thus could provide us with an idea of infection progress. Furthermore, the successfulness of the proposed method in detecting BSR at the very early stage (no symptoms) could prevent the disease widespread.




Fig. (4): Average reflectance of healthy (H) and *G. boninense* infected (U) leaves of F1.

It is also very useful in the real application at the nursery level to test products and screen for tolerant materials.

Conclusion

The differences between healthy and *G. boninense* infected oil palm seedlings were clearly observed in the NIR spectrum where the healthy young leaves gave higher reflectance compared to *G. boninense* infected leaves, although there were no physical symptoms developed. Furthermore, the reflectance at the green spectrum was slightly higher for the healthy leaves. Therefore, it is verified that the usage of F1 was feasible to detect the *G. boninense* disease. This information is useful because this approach can be expended to the more practical application using unmanned aerial vehicle (UAV), where the identification of F1 is easier since it is located at the top of oil palm crown.

Acknowledgement

The authors would like to acknowledge the Ministry of Higher Education Malaysia and Universiti Putra Malaysia for sponsoring this research under Long Term Research Grant

Scheme (LRGS)-Nanomite, research number UPM/700-2/1/LRGS-NANOMITE/5526305.

Conflicts of interest

The authors declare that they have no conflict of interests.

Reference

- Ahmadi, P., Muharam, F. M., Ahmad, K., Mansor, S., & Abu Seman, I. (2017). Early detection of Ganoderma basal stem rot of oil palms using artificial neural network spectral analysis. *Plant Disease*, *101*, 1009-1016. <https://doi.org/10.1094/PDIS-12-16-1699-RE>
- Azuan, N. H., Khairunniza-Bejo, S., Abdullah, A. F., Kassim, M. S. M., & Ahmad, D. (2019). Analysis of changes in oil palm canopy architecture from basal stem rot using terrestrial laser scanner. *Plant Disease*, *103*, 3218-3225. <https://doi.org/10.1094/PDIS-10-18-1721-RE>
- Bravo, C., Moshou, D., West, J., McCartney, A., & Ramon, H. (2003). Early disease detection in wheat fields using spectral reflectance. *Biosystems Engineering*, *84*, 137-145. [https://doi.org/10.1016/s1537-5110\(02\)00269-6](https://doi.org/10.1016/s1537-5110(02)00269-6)
- Chang, C. J. (1998). Pathogenicity of aster yellows phytoplasma and *Spiroplasma citri* on periwinkle. *Phytopathology*, *88*, 1347-1350. <https://doi.org/10.1094/PHYTO.1998.88.12.1347>

- Chappelle, E. W., Kim, M. S., & McMurtrey III, J. E. (1992). Ratio analysis of reflectance spectra (RARS): an algorithm for the remote estimation of the concentrations of chlorophyll a, chlorophyll b, and carotenoids in soybean leaves. *Remote Sensing of Environment*, 39, 239-247. [https://doi.org/10.1016/0034-4257\(92\)90089-3](https://doi.org/10.1016/0034-4257(92)90089-3)
- Darus, A. & Abu Seman, I. (1993). The Ganoderma selective medium (GSM). In *PORIM International Palm Oil Conference. Progress, Prospects Challenges Towards the 21st Century. (Agriculture) September 9-14 Kuala Lumpur, Malaysia* (No. L-0218). PORIM. <http://palmoilis.mpob.gov.my/images/PORIM%20IS/0008/PORIM%20IS%200008.pdf>
- Gamon, J. A., & Surfus, J. S. (1999). Assessing leaf pigment content and activity with a reflectometer. *The New Phytologist*, 143, 105-117. <https://doi.org/10.1046/j.1469-8137.1999.00424.x>
- Gausman, H. W. (1977). Reflectance of leaf components. *Remote Sensing of Environment*, 6, 1-9. [https://doi.org/10.1016/0034-4257\(77\)90015-3](https://doi.org/10.1016/0034-4257(77)90015-3)
- Gitelson, A. A., Merzlyak, M. N., & Chivkunova, O. B. (2001). Optical properties and nondestructive estimation of anthocyanin content in plant leaves. *Photochemistry and Photobiology*, 74, 38-45. [https://doi.org/10.1562/0031-8655\(2001\)074<0038:opaneo>2.0.co;2](https://doi.org/10.1562/0031-8655(2001)074<0038:opaneo>2.0.co;2)
- Gitelson, A. A., Zur, Y., Chivkunova, O. B., & Merzlyak, M. N. (2002). Assessing carotenoid content in plant leaves with reflectance spectroscopy. *Photochemistry and Photobiology*, 75, 272-281. [https://doi.org/10.1562/0031-8655\(2002\)0750272ACIPL2.0.CO2](https://doi.org/10.1562/0031-8655(2002)0750272ACIPL2.0.CO2)
- Husin, N. A., Khairunniza-Bejo, S., Abdullhah, A. F., Kassim, M. S. M., Ahmad, D., & Aziz, M. H. A. (2020). Classification of basal stem rot disease in oil palm plantations using terrestrial laser scanning data and machine learning. *Agronomy* 10, 1624. <https://doi.org/10.3390/agronomy10111624>
- Idris, A. S., & Rafidah, R. (2008). Polyclonal antibody for detection of *Ganoderma*. *MPOB Information Series*, 405. <http://palmoilis.mpob.gov.my/publications/TOT/TT-405.pdf>
- Idris, A. S., Kushairi, D., Ariffin, D., & Basri, M. W. (2006). Technique for inoculation of oil palm germinated seeds with *Ganoderma*. *MPOB Information Series*, 314, 1-4. <http://palmoilis.mpob.gov.my/publications/TOT/TT-314.pdf>
- Idris, A. S., Mazliham, M. S., Loonis, P., & Wahid, M. B. (2010). GanoSken for early detection of *Ganoderma* infection in oil palm. *MPOB Information Series*, 442. <http://palmoilis.mpob.gov.my/publications/TOT/TT-442.pdf>
- Idris, A.S., Yamaoka, M., Hayakawa, S., Basri, M. W., Noorhasimah, I., & Ariffin, D., (2003). PCR technique for detection of *Ganoderma*. *MPOB Information Series*, 188. <http://palmoilis.mpob.gov.my/publications/TOT/tt188.pdf>
- Izzuddin, M. A., Idris, A. S., Nisfariza, N. M., & Ezzati, B. (2015). Spectral based analysis of airborne hyperspectral remote sensing image for detection of *ganoderma* disease in oil palm. In *Proceedings of Conference on Biological and Environmental Science (BIOES 2015)*, 13-20. <https://www.semanticscholar.org/paper/Spectral-based-Analysis-of-Airborne-Hyperspectral-Izzuddin-Idris/827c5439cdeb67d9c4654766cd06a76b9f90e4ee>
- Izzuddin, M. A., Idris, A. S., Wahid, O., Nishfariza, M. N., & Shafri, H. Z. M. (2013). Field spectroscopy for detection of *Ganoderma* disease in oil palm. *MPOB Information Series*, 532. <http://palmoilis.mpob.gov.my/publications/TOT/TT532.pdf>
- Izzuddin, M. A., Idris, A. S., Nisfariza, M. N., Nordiana, A. A., Shafri, H. Z. M., & Ezzati, B. (2017). The development of spectral indices for early detection of *Ganoderma* disease in oil palm seedlings. *International Journal of Remote Sensing*, 38, 6505-6527. <https://doi.org/10.1080/01431161.2017.1335908>
- Kamil, N. N., & Omar, S. F. (2016). Climate variability and its impact on the palm oil industry. *Oil Palm Industry Economic Journal*, 16, 18-30. <http://palmoilis.mpob.gov.my/publications/OPIEJ/opiejv16n1-nadia.pdf>
- Khairunniza-Bejo, S., & Vong, C. N. (2014). Detection of basal stem rot (BSR) infected oil palm tree using laser scanning data. *Agriculture and Agricultural Science Procedia*, 2, 156-164. <https://doi.org/10.1016/J.AASPRO.2014.11.023>

- Kumar, A., Lee, W. S., Ehsani, R. J., Albrigo, L. G., Yang, C., Mangan, R. L. (2012). Citrus greening disease detection using aerial hyperspectral and multispectral imaging techniques. *Journal Applied Remote Sensing*, 6, 063542. <https://doi.org/10.1117/1.JRS.6.063542>
- Liaghat, S., Mansor, S., Ehsani, R., Shafri, H. Z. M., Meon, S., & Sankaran, S. (2014b). Mid-infrared spectroscopy for early detection of basal stem rot disease in oil palm. *Computers and Electronics in Agriculture*, 101, 48-54. <https://doi.org/10.1016/j.compag.2013.12.012>
- Liaghat, S., Ehsani, R., Mansor, S., Shafri, H. Z., Meon, S., Sankaran, S., & Azam, S. H. (2014a). Early detection of basal stem rot disease (*Ganoderma*) in oil palms based on hyperspectral reflectance data using pattern recognition algorithms. *International Journal of Remote Sensing*, 35, 3427-3439. <https://doi.org/10.1080/01431161.2014.903353>
- Lu, J., Ehsani, R., Shi, Y., de Castro, A. I., & Wang, S. (2018). Detection of multi-tomato leaf diseases (late blight, target and bacterial spots) in different stages by using a spectral-based sensor. *Scientific Reports*, 8, 1-11. <https://doi.org/10.1038/s41598-018-21191-6>
- Malenovský, Z., Ufer, C., Lhotakova, Z., Clevers, J. G. P. W., Schaepman, M. E., Albrechtova, J., & Cudlín, P. (2006). A new hyperspectral index for chlorophyll estimation of a forest canopy: Area under curve normalised to maximal band depth between 650-725 nm. *EARSel eProceedings*, 5, 161-172. <https://edepot.wur.nl/39655>
- Moshou, D., Bravo, C., Oberti, R., West, J., Bodria, L., McCartney, A., & Ramon, H. (2005). Plant disease detection based on data fusion of hyper-spectral and multi-spectral fluorescence imaging using Kohonen maps. *Real-Time Imaging*, 11, 75-83. <https://doi.org/10.1016/j.rti.2005.03.003>
- Naidu, Y., Siddiqui, Y., Raffi, M. Y., Saud, H. M., & Idris, A. S. (2018). Inoculation of oil palm seedlings in Malaysia with white-rot hymenomycetes: Assessment of pathogenicity and vegetative growth. *Crop Protection*, 110, 146-154. <https://doi.org/10.1016/j.cropro.2018.02.018>
- Paterson, R. R. M. (2007). *Ganoderma* disease of oil palm—A white rot perspective necessary for integrated control. *Crop protection*, 26, 1369-1376. <https://doi.org/10.1016/j.cropro.2006.11.009>
- Rees, R. W., Flood, J., Hasan, Y., Potter, U., & Cooper, R. M. (2009). Basal stem rot of oil palm (*Elaeis guineensis*); mode of root infection and lower stem invasion by *Ganoderma boninense*. *Plant Pathology*, 58, 982-989. <https://doi.org/10.1111/j.1365-3059.2009.02100.x>
- Sanderson, F. R. (2005). An insight into spore dispersal of *Ganoderma boninense* on oil palm. *Mycopathologia*, 159, 139-141. <https://doi.org/10.1007/s11046-004-4436-2>
- Sariah, M., Hussin, M. Z., Miller, R. N. G., & Holderness, M. (1994). Pathogenicity of *Ganoderma boninense* tested by inoculation of oil palm seedlings. *Plant Pathology*, 43, 507-510. <https://doi.org/10.1111/j.1365-3059.1994.tb01584.x>
- <https://doi.org/10.1080/01431161.2010.519003>
- Shafri, H. Z. M., & Hamdan, N. (2009). Hyperspectral imagery for mapping disease infection in oil palm plantation using vegetation indices and red edge techniques. *American Journal of Applied Sciences*, 6, 1031-1035. <https://doi.org/10.3844/ajassp.2009.1031.1035>
- Shafri, H. Z. M., Hamdan, N., & Izzuddin Anuar, M. (2012). Detection of stressed oil palms from an airborne sensor using optimised spectral indices. *International Journal of Remote Sensing*, 33, 4293-4311. <https://doi.org/10.1080/01431161.2011.619208>
- Shafri, H. Z., Anuar, M. I., Seman, I. A., & Noor, N. M. (2011). Spectral discrimination of healthy and *Ganoderma*-infected oil palms from hyperspectral data. *International Journal of Remote Sensing*, 32, 7111-7129.
- Schmidt, K. S., & Skidmore, A. K. (2003). Spectral discrimination of vegetation types in a coastal wetland. *Remote Sensing of Environment*, 85, 92-108. [https://doi.org/10.1016/S0034-4257\(02\)00196-7](https://doi.org/10.1016/S0034-4257(02)00196-7)
- Slaton, M. R., Raymond Hunt Jr, E., & Smith, W. K. (2001). Estimating near-infrared leaf reflectance from leaf structural characteristics. *American Journal of Botany*, 88, 278-284. <https://doi.org/10.2307/2657019>
- Shu'ud, M. M., Loonis, P., & Seman, I. A. (2007). Towards automatic recognition and grading of *Ganoderma* infection pattern using fuzzy systems. *International Journal of Medical Health Sciences*, 1, 1-6. <https://publications.waset.org/5920/towards->

automatic-recognition-and-grading-of-ganoderma-infection-pattern-using-fuzzy-systems

Susič, N., Žibrat, U., Širca, S., Strajnar, P., Razinger, J., Knapič, M., & Stare, B. G. (2018). Discrimination between abiotic and biotic drought stress in tomatoes using hyperspectral imaging. *Sensors and Actuators B: Chemical*, 273, 842-852. <https://doi.org/10.1016/j.snb.2018.06.121>

Zarco-Tejada, P. J., Miller, J. R., Morales, A., Berjón, A., & Agüera, J. (2004). Hyperspectral indices and

model simulation for chlorophyll estimation in open-canopy tree crops. *Remote Sensing of Environment*, 90, 463-476. <https://doi.org/10.1016/j.rse.2004.01.017>

Zwiggelaar, R. (1998). A review of spectral properties of plants and their potential use for crop/weed discrimination in row-crops. *Crop Protection*, 17, 189-206. [https://doi.org/10.1016/S0261-2194\(98\)00009-X](https://doi.org/10.1016/S0261-2194(98)00009-X)